

Product Name: SAME

Primary Ingredient(s): S-Adenosyl-L-methionine (SAME)

References S-Adenosyl-L-methionine (SAME):

- Bressa GM. S-Adenosyl-L-methionine (SAME) as antidepressant: meta-analysis of clinical studies. *Acta Neurol Scand* 1994;154:7–14.
- Lieber CS. Role of S-adenosyl-L-methionine in the treatment of liver diseases. *J Hepatol* 1999;30:1155–9.
- Williams R, Lieber CS. The role of SAME in the treatment of liver disease. *Drugs* 1990;40(suppl):1–2.
- Mato JM, Camara J, Fernandez de Paz J, et al. S-Adenosylmethionine in alcoholic liver cirrhosis: a randomized, placebo-controlled, double-blind, multicenter clinical trial. *J Hepatol* 1999;30:1081–9.
- Brown R, Gerbarg P, Bottiglieri T. S-Adenosylmethionine in the clinical practice of psychiatry, neurology and internal medicine. *Clin Pract Intern Med* 2000;1:230–41.
- Algeri S, Catto E, Curcio M, Ponzio F, Stramentinoli G. Changes in rat brain noradrenaline and serotonin after administration of S-adenosylmethionine. In: Zappia V, Usdin E, Salvatore S, eds. *Biochemical and pharmacological roles of adenosylmethionine and the central nervous system*. New York: Pergamon Press, 1979:81–7.
- Otero-Losado ME, Rubio MC. Acute effects of S-adenosyl-L-methionine on catecholaminergic central function. *Eur J Pharmacol* 1989;163:353–6.
- Otero-Losado ME, Rubio MC. Acute changes in 5HT metabolism after S-adenosylmethionine administration. *Gen Pharmacol* 1989;20:403–6.
- Curcio M, Catto E, Stramentinoli G, Algeri S. Effect of SAME on 5HT metabolism in rat brain. *Prog Neuropsychopharmacol* 1978;2:65–71.
- Loguercio C, Nardi G, Argenzio F, et al. Effect of S-adenosyl-L-methionine administration on red blood cell cysteine and glutathione levels in alcoholic patients with and without liver disease. *Alcohol Alcohol* 1994;29:597–604.

-
- Galan AI, Munoz ME, Palomero J, Moreno C, Jimenez R. Role of S-adenosylmethionine on the hepatobiliary homeostasis of glutathione during cyclosporine A treatment. *J Physiol Biochem* 2000;56:189–200.
 - Vendemiale G, Altomare E, Trizio T, et al. Effects of oral S-adenosyl-L-methionine on hepatic glutathione in patients with liver disease. *Scand J Gastroenterol* 1989;24:407–15.
 - Lieber CS. Alcoholic liver disease: new insights in pathogenesis lead to new treatments. *J Hepatol* 2000;32(suppl):113–28.
 - Mato JM, Alvarez L, Ortiz P, Pajares MA. S-Adenosylmethionine synthesis: molecular mechanisms and clinical implications. *Pharmacol Ther* 1997;73:265–80.
 - Agnoli A, Andreoli V, Casacchia M, Cerbo R. Effect of S-adenosyl-L-methionine (SAMe) upon depressive symptoms. *J Psychiatr Res* 1976;13:43–54.
 - Di Padova C. S-Adenosylmethionine in the treatment of osteoarthritis. Review of the clinical studies. *Am J Med* 1987;83:60–5.
 - Harmand MF, Vilamitjana J, Maloche E, Duphil R, Ducassou D. Effects of S-adenosylmethionine on human articular chondrocyte differentiation. An in vitro study. *Am J Med* 1987;83:48–54.
 - Barcelo HA, Wiemeyer JC, Sagasta CL, Macias M, Barreira JC. Effect of S-adenosylmethionine on experimental osteoarthritis in rabbits. *Am J Med* 1987;83:55–9.
 - Gutierrez S, Palacios I, Sanchez-Pernaute O, et al. SAMe restores the changes in the proliferation and in the synthesis of fibronectin and proteoglycans induced by tumour necrosis factor alpha on cultured rabbit synovial cells. *Br J Rheumatol* 1997;36:27–31.
 - Bradley JD, Flusser D, Katz BP, et al. A randomized, double blind, placebo controlled trial of intravenous loading with S-adenosylmethionine (SAM) followed by oral SAM therapy in patients with knee osteoarthritis. *J Rheumatol* 1994;21:905–11.